



دانشگاه علوم پزشکی قزوین

دانشکده پرستاری و مامایی

پایان نامه برای دریافت درجه

کارشناسی ارشد رشته پرستاری ویژه

عنوان پایان نامه

بررسی ارتباط همدلی و نعدوستی با نگرش پرستاران بخش های ICU، اورژانس و دیالیز در خصوص اهدا عضو در مراکز آموزشی درمانی وابسته به دانشگاه علوم پزشکی قزوین در سال ۱۳۹۸

استاد راهنما:

سرکار خانم دکتر یکه فلاح

استاد مشاور:

سرکار خانم لیلا دهقانکار

استاد مشاور آمار:

آقای دکتر مهدی رنجبران

نگارنده:

مهناز طاهرخانی

ماه و سال انتشار:

بهمن ماه سال ۱۳۹۸

بررسی ارتباط همدلی و نوع و دیالیز در خصوص اهدای عضو در مراکز آموزشی درمانی وابسته به دانشگاه علوم پزشکی

قزوین در سال ۱۳۹۸

چکیده:

زمینه و هدف: پرستاران بخش های مراقبت های ویژه نقش مهمی در روند اهدای عضو دارند و عملکرد پرستاران در رابطه با ایفای نقش خود در فرآیند اهدای عضو بیشتر متاثر از نگرش آنها در این رابطه است. لذا شناخت عوامل موثر بر نگرش مثبت پرستاران به اهدای عضو امری ضروری است. این مطالعه با هدف تعیین ارتباط بین همدلی و نودوستی با نگرش پرستاران بخش های ویژه نسبت به اهدای عضو و مرگ مغزی در قزوین انجام گردید.

مواد و روش ها : در این مطالعه توصیفی مقطعی که در سال ۱۳۹۸ در بیمارستانهای آموزشی درمانی دانشگاه علوم پزشکی قزوین صورت گرفت تعداد ۲۲۲ نفر از پرستاران بخش های ویژه به روش نمونه گیری تصادفی طبقه ای وارد مطالعه شدند. برای گردآوری داده ها پرسشنامه های استاندارد نودوستی ، همدلی و نگرش به اهدای عضو توسط پرستاران تکمیل گردید.

یافته ها: بین نودوستی ($p < 0.001$ و $r = 0.24$) و مولفه های آن: رفتارهای اجتماعی مطلوب ناشناس (گمنام) ($p < 0.001$) و رفتارهای اجتماعی مطلوب هیجانی ($p = 0.03$ و $r = 0.14$) ، رفتارهای اجتماعی مطلوب در موقعیت اضطراری ($p = 0.03$ و $r = 0.14$) ، رفتارهای اجتماعی مطلوب متابعت آمیز ($p = 0.001$ و $r = 0.21$) با نگرش پرستاران نسبت به اهدا عضو ، همبستگی مثبت و معنی دار وجود داشت. همچنین ارتباط بین همدلی ($p = 0.04$ و $r = 0.13$) و مولفه دیدگاه گیری ($p = 0.02$ و $r = 0.152$) و همدلی خیالی ($p = 0.03$ و $r = 0.14$) با نگرش پرستاران ، از لحاظ آماری معنی دار بود. جهت بررسی پیش بینی کننده های نگرش به اهدا عضو ، متغیرهایی که در مدل تک متغیره دارای $p < 0.05$ بودند وارد رگرسیون خطی چندگانه به روش inter شدند که طبق نتایج مولفه ی رفتارهای اجتماعی مطلوب ناشناس (گمنام) از متغیر نودوستی ($p < 0.0001$) و متغیرهای جنس ($p = 0.007$) و داشتن کارت اهدا عضو ($p = 0.012$) ، به عنوان پیش بینی کننده های مثبت نگرش پرستاران در مورد اهدا عضو بودند و در مجموع این متغیرها توانستند ۲۰٪ تغییرات نگرش را پیش بینی کنند.

نتیجه گیری: نتایج نشان داد که نوع دوستی و همدلی از مهمترین عوامل تمایل پرستاران به اهدای عضو است. بنابراین ، اجرای برنامه های آموزشی در رابطه با تغییر نگرش پرستار به منظور افزایش احساس نودوستی و همدلی در مورد اهدای عضو ضروری است

کلید واژه ها: نودوستی، همدلی، نگرش، اهدا عضو، پرستاران



**Qazvin University of Medical Sciences
School of Nursing & Midwifery**

**Thesis submitted to achieve the Degree of M.Sc.
In Critical Care Nursing**

Title

The association between Empathy and Altruism with nurses' attitudes towards organ donation in ICU, Dialysis and Emergency departments in clinical-educational centers affiliated to Qazvin University of Medical Sciences in 2019

Under supervision of:

Leili yekehfallah

Advisers:

Leila dehghankar

Mehdi ranjbaran

By:

Mahnaz taherkhani

Month & year

January 2020

The association between Empathy and Altruism with nurses' attitudes towards organ donation in ICU, Dialysis and Emergency departments in clinical-educational centers affiliated to Qazvin University of Medical Sciences in 2019

Abstract:

Aims:

Nurses, especially nurses in ICUs, play an important role in organ donation. Nurses performance in relation to their roles is more affected by their attitudes in this regard, therefore identification of effective factors on the positive attitudes of the nurses towards organ donation is essential. This study aimed to determine the association between empathy and altruism with the nurses' attitudes in intensive care unit towards organ donation and brain death.

Methods:

In this cross-sectional study, which was conducted in 2019, 222 nurses from intensive care units were selected by stratified random sampling. For collecting data, the questionnaires of altruism, empathy and attitude toward organ donation were completed by nurses. Data was analyzed using Pearson correlation and multiple linear regression.

Result:

There was a positive and significant correlation between altruism ($p < 0.001$, $r = 0.24$) and its components (anonymous prosocial behaviors ($r = 0.33$, $p < 0.001$), emotional prosocial behaviors ($r = 0.14$, $p = 0.03$), dire prosocial behaviors ($r = 0.14$, $p = 0.03$) and compliant prosocial behaviors ($r = 0.21$ and $p = 0.001$)) with nurses' attitudes. Also the association between empathy ($r = 0.04$, $r = 0.13$), perspective taking component ($p = 0.02$, $r = 0.152$) and imaginary empathy ($r = 0.14$, $p = 0.03$) with nurses' attitudes was statistically significant. The components of anonymous prosocial behaviors ($p < 0.0001$), gender ($p = 0.007$) and having organ donation card ($p = 0.012$) were positive predictors of nurses' attitudes towards organ donation. To investigate the predictors of attitudes toward organ donation, variables in univariate model with $p < 0.2$ were entered into multiple linear regression by inter method. According to the results, the component of anonymous prosocial behaviors in altruism ($p < 0.0001$), gender ($p = 0.007$) and having organ donation card ($p = 0.012$) were positive predictors of nurses' attitude toward organ donation, these variables were able to predict 20% of attitude changes.

Conclusion:

The results showed that altruism and empathy were most important factors in the tendency of nurses to organ donation. Therefore, the implementation of educational programs regarding changing nurse's attitude in order to increase the sense of altruism and empathy about organ donation is necessary.

Key words: Altruism, Empathy, Attitude, Organ Donation, Nurses